arXiv:math/0103193v2 [math.CT] 28 Sep 2009

On Ext in the Category of Functors to Preabelian Category *

A. A. Husainov
Komsomolsk-on-Amur State Technical University,
Faculty of Computer Technologies,
681013, Komsomolsk-on-Amur, prosp. Lenina, 27, Russia
husainov@knastu.ru

A. Pancar
Ondokuz Mayis University, Faculty of Art and Sciences,
Department of Mathematics, 55139, Kurupelit, Samsun, Turkey
apancar@omu.edu.tr

M. Yapici
Ondokuz Mayis University, Faculty of Art and Sciences,
Department of Mathematics, 55139, Kurupelit, Samsun, Turkey
myapici@omu.edu.tr

1991 AMS Subject Classification: 18G25

Abstract

The article is devoted to extension groups in the category of functors from a small category to an additive category with an Abelian structure in the sense of Heller. It is proved that under additional assumptions there exists a spectral sequence which converges to the extension groups.

Key words : *Extension groups, cohomology of categories.*

1 INTRODUCTION

Let Ab be the category of abelian groups, Mod_R the category of left R-modules over a ring with identity.

Jibladze and Pirashvili [4], [5] proved that if $\mathcal{A} = \mathrm{Mod}_R$ then for every small category $\mathbb{C}$ and functors $F, G : \mathbb{C} \to \mathcal{A}$ there exists a first quadrant spectral sequence with

$$E_2^{p,q} = H^p(\mathbb{C}, \{\mathrm{Ext}^q(F(a), G(b))\})$$

---

*In the preparation of this paper, we have been assisted by a grant from the TÜBİTAK and NATO. The first author is supported by Russian Federation Education Department

which converges to $\mathrm{Ext}^n(F,G)$. Here $H^n(\mathbb{C},B)$ are the n-th Hochschild-Mitchell cohomology groups with coefficients in a bimodule $B : \mathbb{C}^{op}\times\mathbb{C} \to \mathrm{Ab}$ in the sense of [1], and $\{\mathrm{Ext}^q(F(a),G(b))\}$ is the functor $\mathrm{Ext}^q(F(-),G(=)) : \mathbb{C}^{op}\times\mathbb{C} \to \mathrm{Ab}$. This spectral sequence was generalized by Khusainov [6] to an Abelian category $\mathcal{A}$ with a proper class [9]. We generalize it to additive categories with Abelian structures in the sense of Heller [3].

2 ABELIAN STRUCTURE

Let $\mathcal{A}$ be a preadditive category, $B \in \mathcal{A}$ an object. A *direct sum decomposition* (A, A', i, p, i', p') of B consist of objects and morphisms

$$A \xrightarrow{i} B \xleftarrow{i'} A', \quad A \xleftarrow{p} B \xrightarrow{p'} A'$$

satisfying to the conditions:

$$\begin{gathered} p \circ i = 1_A, \quad p' \circ i' = 1_{A'}, \\ p' \circ i = 0_{AA'}, \quad p \circ i' = 0_{A'A}, \\ i \circ p + i' \circ p' = 1_B \end{gathered}$$

A morphism $\rho : B \to A$ is called a *retraction* if there exists a morphism $\nu : A \to B$ such that $\rho \circ \nu = 1_A$.

An *additive* category is a preadditive category with a zero-object and with finite products. We follow to Mac Lane [9] for the notions of the kernels and cokernels.

Lemma 2.1 *Let $\mathcal{A}$ be an additive category. Then the following conditions are equivalent:*

(1) Every retraction has a kernel;

(2) For every morphisms $\rho : B \to A$ and $i : A \to B$ in $\mathcal{A}$ satisfying $\rho \circ i = 1_A$ there are an object A' and morphisms $\rho' : B \to A'$, $i' : A' \to B$ such that $(A, A', i, \rho, i', \rho')$ is a direct sum decomposition of B.

The condition (2) is called Cancellation Axiom in [3]. Thus, if $\mathcal{A}$ is an additive category in which all retractions have kernels then $\mathcal{A}$ is a category with Cancellation Axiom.

The sequence of morphisms

$$0 \longrightarrow A' \xrightarrow{\alpha'} A \xrightarrow{\alpha''} A'' \longrightarrow 0$$

is called a *short exact sequence*, abbreviated "s.e.s", if (A'', α'') is a cokernel of α' and (A', α') is a kernel of α''. More generally, a sequence

$$A_n \xrightarrow{f_n} A_{n-1} \xrightarrow{f_{n-1}} \dots \xrightarrow{f_1} A_0$$

is *exact* if there are s.e.s $0 \to B_j \overset{u_j}{\to} A_j \overset{\nu_j}{\to} B_{j-1} \to 0$ for $j = 1, 2, \ldots, n-1$, an epimorphism $\nu_n : A_n \to B_{n-1}$ and a monomorphism $u_0 : B_0 \to A_0$ such that $f_j = u_{j-1} \circ \nu_j, \quad j = 1, 2, \ldots, n.$

Let $\mathcal{A}$ be an additive category with Cancellation Axiom. An *Abelian structure* on $\mathcal{A}$ is a subclass $\mathcal{P} \subseteq \mathrm{Mor}\mathcal{A}$ whose elements are called to be *proper maps.* A short exact sequence whose maps are proper is a *proper s.e.s.*

The class $\mathcal{P}$ is to be subject to the following axioms:

(P0) $1_A \in \mathcal{P}$ for all $A \in \mathrm{Ob}\mathcal{A}$.

(P1) If $(f : B \to C) \in \mathcal{P}$, $(g : A \to B) \in \mathcal{P}$ and g is an epimorphism then $f \circ g \in \mathcal{P}$; dually, if $f \in \mathcal{P}$, $g \in \mathcal{P}$ and f is a monomorphism then $f \circ g \in \mathcal{P}$.

(P2) If $f \circ g \in \mathcal{P}$ is a monomorphism then g is proper; dually, if $f \circ g \in \mathcal{P}$ is an epimorphism then f is proper.

(P3) For every proper map $f : B \to D$ there are proper s.e.s $0 \to A \to B \to C \to 0$ and $0 \to C \to D \to E \to 0$ such that the following diagram

$$\begin{tikzcd}
0 \arrow[r] & A \arrow[r] & B \arrow[r] \arrow[d, "f"] & C \arrow[r] \arrow[d, "1_C"] & 0 \\
0 & E \arrow[l] & D \arrow[l] & C \arrow[l] & 0 \arrow[l]
\end{tikzcd}$$

is commutative.

(P4) If in the commutative diagram

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & A' \arrow[r] \arrow[d] & A \arrow[r] \arrow[d] & A'' \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & B' \arrow[r] \arrow[d] & B \arrow[r] \arrow[d] & B'' \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & C' \arrow[r] \arrow[d] & C \arrow[r] \arrow[d] & C'' \arrow[r] \arrow[d] & 0 \\
 & 0 & 0 & 0 &
\end{tikzcd}$$

all collumns and the second two rows are proper s.e.s, then the first row is a proper s.e.s.

We remark that (P3) implies the existence of kernels and cokernels for all proper maps by [3, Prop. 2.2].

Let $\mathcal{A}$ be an additive category with Cancellation Axiom, $\mathbb{C}$ a small category. We denote by $\mathcal{A}^{\mathbb{C}}$ the category of functors $\mathbb{C} \to \mathcal{A}$. If $\rho : F \to G$ is a retraction in $\mathcal{A}^{\mathbb{C}}$ then $\rho_c : F(c) \to G(c)$ are retractions for all $c \in \mathrm{Ob}\mathbb{C}$ and have kernels by Lemma 2.1. Consequently, $\rho : F \to G$ has a kernel. Therefore $\mathcal{A}^{\mathbb{C}}$ is the additive category with Cancellation Axiom.

The Abelian structure generalizes the proper class in an Abelian category in the sense of [9]. We refer to [7] and [3] for other examples of Abelian structure.

By [3, Prop. 3.5] an additive category carries Abelian structures if and only if it satisfies the Cancellation Axiom.

Proposition 2.2 *Let $\mathcal{A}$ be an additive category with an Abelian structure $\mathcal{P}$, $\mathbb{C}$ a small category, $\mathbb{C}\mathcal{P}$ the class of all natural transformations $\eta : F \to G$ for which $\eta_c \in \mathcal{P}$ for all $c \in \mathrm{Ob}\mathbb{C}$. Then the class $\mathbb{C}\mathcal{P}$ is an Abelian structure on $\mathcal{A}^{\mathbb{C}}$.*

Proof. It is clear that $\mathcal{A}^{\mathbb{C}}$ is the category with Cancellation Axiom by Lemma 2.1. We check the axioms (P0)-(P4) for $\mathbb{C}\mathcal{P}$:

(P0) Each identity $1_F : F \to F$ consists of $1_{F(c)}$, $c \in \mathrm{Ob}\mathbb{C}$. Hence $1_F \in \mathcal{P}$.

(P1) A morphism $\eta : F \to G$ in $\mathcal{A}^{\mathbb{C}}$ is a monomorphism if and only if $\ker \eta = 0$. Hence η is a monomorphism if and only if η_c are monomorphisms for all $c \in \mathrm{Ob}\mathbb{C}$; dually for epimorphisms.

(P2) Analogously.

(P3) For every $f \in \mathcal{P}$ we choose a kernel $(\ker f, k_f)$ of f. By definition of kernels [9, Ch.IX] for every commutative diagram

$$\begin{tikzcd} A \arrow[r, "f"] \arrow[d, "\alpha"'] & B \arrow[d, "\beta"] \\ A' \arrow[r, "f'"] & B' \end{tikzcd} \tag{1}$$

with $f, f' \in \mathcal{P}$ and $\alpha, \beta \in \mathrm{Mor}\mathcal{A}$ there exists unique $\ker f \to \ker f'$ for which the following diagram is commutative

$$\begin{tikzcd} \ker f \arrow[r, "k_f"] \arrow[d] & A \arrow[r, "f"] \arrow[d, "\alpha"] & B \arrow[d, "\beta"] \\ \ker f' \arrow[r, "k_{f'}"] & A' \arrow[r, "f'"] & B' \end{tikzcd}$$

Dually, we choose cokernels $(\mathrm{coker} f, c_f)$. Consider the category which objects are all members of $\mathcal{P}$ and morphisms are commutative diagrams (1).

We see that ker and coker are functors from this category to $\mathcal{A}$. Hence for every natural transformation $\eta : F \to G$ with $\eta_c \in \mathcal{P}$, for all $c \in \mathrm{Ob}\mathbb{C}$, we have a diagram of natural transformations

$$\begin{tikzcd}
0 \arrow[r] & \ker\eta \arrow[r, "k_\eta"] & F \arrow[r, "\varepsilon"] \arrow[d, "\eta"] & \mathrm{coker} k_\eta \arrow[r] \arrow[d, "\xi"] & 0 \\
0 & \mathrm{coker}\eta \arrow[l] & G \arrow[l, "c_\eta"'] & \ker c_\eta \arrow[l, "m"'] & 0 \arrow[l]
\end{tikzcd}$$

where ξ is an isomorphism of functors. By [3, Prop.3.3.] all equivalences in $\mathcal{A}$ belong to $\mathcal{P}$, therefore $\xi \circ \varepsilon$ is a proper epimorphism, and we obtain (P3) for $\mathbb{C}\mathcal{P}$.

(P4) *A* sequence of functors is exact if and only if it is exact at every $c \in \mathrm{Ob}\mathbb{C}$. Thus (P4) holds.

Q.E.D.

An object P is called $\mathcal{P}$-projective if for every proper epimorphism $\varepsilon : A \to B$ and for a morphism $\alpha : P \to B$ there is $\beta : P \to A$ such that $\varepsilon \circ \beta = \alpha$.

Let $\mathcal{A}$ be an additive category with Cancellation Axiom, $\mathcal{P} \subseteq \mathrm{Mor}\mathcal{A}$ an Abelian structure. Following Heller we will say that $\mathcal{A}$ has enough $\mathcal{P}$-projectives if for each $A \in \mathrm{Ob}\mathcal{A}$ there is a $\mathcal{P}$-projective object $P(A) \in \mathcal{A}$ and a proper epimorphism $P(A) \overset{\pi A}{\to} A$.

Proposition 2.3 *Let $\mathcal{A}$ be an additive category with coproducts, $\mathcal{P}$ an Abelian structure. If $\mathcal{A}$ has enough $\mathcal{P}$-projectives and coproducts of proper epimorphisms are proper, then $\mathcal{A}^{\mathbb{C}}$ has enough $\mathbb{C}\mathcal{P}$-projectives.*

PROOF. We consider the set $\mathrm{Ob}\mathbb{C}$ as the maximal discrete subcategory of $\mathbb{C}$. Let $O : \mathcal{A}^{\mathbb{C}} \to \mathcal{A}^{\mathrm{Ob}\mathbb{C}}$ be the restriction functor, $O(F) = F|_{\mathrm{Ob}\mathbb{C}}$. The category $\mathcal{A}$ has coproducts, hence there is a left adjoint functor Λ to the functor O. Up to a natural isomorphism, we have $\Lambda(D)(c) = \sum_{c_0 \to c} D(c_o)$ for each $D \in \mathcal{A}^{\mathrm{Ob}\mathbb{C}}$. The counit of adjunction $\varepsilon_F : \Lambda O F \to F$ splits on each $c \in \mathrm{Ob}\mathbb{C}$, hence ε_F is a proper epimorphism in $\mathcal{A}^{\mathbb{C}}$ with respect to Abelian structure $\mathbb{C}\mathcal{P}$. Choose a family $P = \{P(c)\}_{c \in \mathrm{Ob}\mathbb{C}} \in \mathcal{A}^{\mathrm{Ob}\mathbb{C}}$ of $\mathcal{P}$-projectives and a family of proper epimorphisms $\psi = \{\psi_c : P(c) \to F(c)\}_{c \in \mathrm{Ob}\mathbb{C}}$ and apply the functor Λ. Then $\Lambda(\psi)_c : \Lambda(P)(c) \to \Lambda(OF)(c)$ is a proper epimorphism for each $c \in \mathrm{Ob}\mathbb{C}$ as the coproduct of proper epimorphisms.

Consequently, $\varepsilon_F \circ (\Lambda(\psi)) : \Lambda(P) \to F$ is a proper epimorphism in $\mathcal{A}^{\mathbb{C}}$.

The functor $\mathcal{A}^{\mathbb{C}}(\Lambda(P), -)$ is isomorphic to the functor $\mathcal{A}^{\mathrm{Ob}\mathbb{C}}(P, O(-))$ which is exact on all proper s.e.s in $\mathcal{A}^{\mathbb{C}}$. Thus, $\Lambda(P)$ is $\mathbb{C}\mathcal{P}$-projective and there is an proper epimorphism $\Lambda(P) \to F$. Q.E.D.

3 COHOMOLOGY OF CATEGORIES

Let $\mathbb{C}$ be a small category, $F : \mathbb{C} \to \mathrm{Ab}$ a functor.

Consider the sequence of groups

$$C^n(\mathbb{C}, F) = \prod_{c_0 \to \ldots \to c_n} F(c_n), \quad n \geq 0.$$

Let $N_n\mathbb{C}$ be the set of all sequences of morphisms $c_0 \to c_1 \to \ldots \to c_n$ in $\mathbb{C}$. Regarding each $\varphi \in \prod_{c_0 \to \ldots \to c_n} F(c_n)$ as a function $\varphi : N_n\mathbb{C} \to \bigcup_{c \in \mathrm{Ob}\mathbb{C}} F(c)$ with $\varphi(c_0 \to \ldots \to c_n) \in F(c_n)$, we define the homomorphisms $d^n : C^n(\mathbb{C}, F) \to C^{n+1}(\mathbb{C}, F)$ by the formulas

$$\begin{gathered}(d^n\varphi)(c_0 \overset{\alpha_1}{\to} c_1 \to \ldots \overset{\alpha_{n+1}}{\to} c_{n+1}) = \\ \sum_{i=0}^{n} (-1)^i \varphi(c_0 \overset{\alpha_1}{\to} \ldots \overset{\alpha_i}{\to} \hat{c}_i \overset{\alpha_{i+1}}{\to} \ldots \overset{\alpha_{n+1}}{\to} c_{n+1}) + \\ (-1)^{n+1} F(\alpha_{n+1})(\varphi(c_0 \overset{\alpha_1}{\to} \ldots \overset{\alpha_n}{\to} c_n)).\end{gathered}$$

Here $(c_0 \overset{\alpha_1}{\to} \ldots \overset{\alpha_i}{\to} \hat{c}_i \overset{\alpha_{i+1}}{\to} \ldots \overset{\alpha_{n+1}}{\to} c_{n+1})$ is equal to $(c_0 \overset{\alpha_1}{\to} \ldots \to c_{i-1} \xrightarrow{\alpha_{i+1} \circ \alpha_i} c_{i+1} \to \ldots \overset{\alpha_{n+1}}{\to} c_{n+1})$ if $0 < i < n+1$, and to $(c_1 \overset{\alpha_2}{\to} \ldots \overset{\alpha_{n+1}}{\to} c_{n+1})$ if $i = 0$. It is well known that $d^{n+1} \circ d^n = 0$ for all $n \geq 0$. We have obtained the complex of Abelian groups and homomorphisms

$$0 \to C^0(\mathbb{C}, F) \overset{d^0}{\to} \ldots \to C^n(\mathbb{C}, F) \overset{d^n}{\to} C^{n+1}(\mathbb{C}, F) \to \ldots$$

which is denoted by $C^*(\mathbb{C}, F)$. The cohomologies $H^n(C^*(\mathbb{C}, F)) = \ker d^n / \mathrm{Im} d^{n-1}$ are isomorphic to Abelian groups $\lim^n_{\mathbb{C}} F$ (see [11], for example) where $\lim^n_{\mathbb{C}} : \mathrm{Ab}^{\mathbb{C}} \to \mathrm{Ab}$ are the n-th right derived functors of the limit $\lim_{\mathbb{C}} : \mathrm{Ab}^{\mathbb{C}} \to \mathrm{Ab}$.

In particular, if $\mathbb{C}$ has the initial object then $H^n(C^*(\mathbb{C}, F)) = 0$ for $n > 0$ and $H^0(C^*(\mathbb{C}, F)) \cong \lim_{\mathbb{C}} F$.

For each $c \in \mathrm{Ob}\mathbb{C}$ we denote by $c/\mathbb{C}$ the *comma-category* in the sense of [8], its objects are pairs (a, α) of $a \in \mathrm{Ob}\mathbb{C}$ and $\alpha \in \mathbb{C}(c, a)$, and for each pair of objects $(a, \alpha \in \mathbb{C}(c, a))$ and $(b, \beta \in \mathbb{C}(c, b))$ the set of morphisms from (a, α) to (b, β) consists of $\gamma \in \mathbb{C}(a, b)$ for which the following diagrams

$$\begin{tikzcd} c \arrow[r, "\alpha"] \arrow[d, "="] & a \arrow[d, "\gamma"] \\ c \arrow[r, "\beta"'] & b \end{tikzcd} \tag{2}$$

are commutative.

Let $Q_c : c/\mathbb{C} \to \mathbb{C}$ be the functor which carries the above diagram (2) to $\gamma : a \to b$. The category $c/\mathbb{C}$ contains the initial object $(c, 1_c)$. Therefore, the functor $\lim_{c/\mathbb{C}}$ is exact and $H^n(C^*(\mathbb{C}, GQ_c)) = 0$ for every functor $F : \mathbb{C} \to \mathrm{Ab}$ and $n > 0$.

Let $\mathbb{C}$ be a small category. For each $a \in \mathrm{Ob}\mathbb{C}$ we identify the morphism 1_a with the object a, so $\mathrm{Ob}\mathbb{C} \subseteq \mathrm{Mor}\mathbb{C}$. Objects of the *factorization category* [1] $\mathbb{C}'$ are all morphisms of $\mathbb{C}$, and the set of morphisms $\mathbb{C}'(f,g)$ for any $f, g \in \mathrm{Mor}\mathbb{C}$ consists of pairs (α, β) of morphisms in $\mathbb{C}$ for which the diagram

$$\begin{tikzcd}
b \arrow[r, "\beta"] & d \\
a \arrow[u, "f"] & c \arrow[l, "\alpha"] \arrow[u, "g"']
\end{tikzcd}$$

is commutative; we denote the morphisms by $(\alpha, \beta) : f \to g$. The composition is defined as $(\alpha_2, \beta_2) \circ (\alpha_1, \beta_1) = (\alpha_1 \circ \alpha_2, \beta_2 \circ \beta_1)$. Functors $F : \mathbb{C}' \to \mathrm{Ab}$ are called *natural systems.*

Let $\mathbb{C}$ be a small category, $F : \mathbb{C}' \to \mathrm{Ab}$ a natural system. For every $n \geq 0$ we consider an Abelian group

$$K^n(\mathbb{C}, F) = \prod_{c_0 \overset{\alpha_1}{\to} c_1 \overset{\alpha_2}{\to} c_2 \to \ldots \overset{\alpha_n}{\to} c_n} F(\alpha_n \circ \alpha_{n-1} \circ \ldots \circ \alpha_1)$$

We regard elements of $K^n(\mathbb{C}, F)$ as maps $\varphi : N_n\mathbb{C} \to \bigcup_{g \in \mathrm{Mor}\mathbb{C}} F(g)$ with $\varphi(\alpha_1, \ldots, \alpha_n) \in F(\alpha_n \circ \alpha_{n-1} \circ \cdots \circ \alpha_1)$ and define homomorphisms $d^n : K^n(\mathbb{C}, F) \to K^{n+1}(\mathbb{C}, F)$ by the formula

$$\begin{gathered}(d^n\varphi)(\alpha_1, \ldots, \alpha_{n+1}) = F(\alpha_1, 1)\varphi(\alpha_2, \ldots, \alpha_{n+1}) + \\ \sum_{i=1}^{n} (-1)^i \varphi(\alpha_1, \ldots \alpha_{i-1}, \alpha_{i+1} \circ \alpha_i, \alpha_{i+2}, \ldots, \alpha_{n+1}) + \\ (-1)^{n+1} F(1, \alpha_{n+1})\varphi(\alpha_1, \ldots, \alpha_n).\end{gathered}$$

Cohomology groups $H^n(K^*(\mathbb{C}, F)) = \ker d^n / \mathrm{Im} d^{n-1}$ are called n-th cohomology groups $H^n(\mathbb{C}, F)$ of $\mathbb{C}$ with coefficients in the natural system F.

For any $\alpha \in \mathrm{Mor}\mathbb{C}$ we denote by $\mathrm{dom}\alpha$ and $\mathrm{cod}\alpha$ the domain and the codomain. We have the functor $(\mathrm{dom}, \mathrm{cod}) : \mathbb{C}' \to \mathbb{C}^{op} \times \mathbb{C}$ which acts as $f \mapsto (\mathrm{dom}f, \mathrm{cod}f)$ on objects, and $((\alpha, \beta) : f \to g) \mapsto (\alpha, \beta)$ on morphisms of $\mathbb{C}'$. Baues and Wirsching [1] proved that $H^n(\mathbb{C}, F) \cong \lim^n_{\mathbb{C}'} F$ for all $n \geq 0$. Functors $B : \mathbb{C}^{op} \times \mathbb{C} \to \mathrm{Ab}$ are called *bimodules* over $\mathbb{C}$. For each bimodule B over $\mathbb{C}$ we denote by $\{B(\mathrm{dom}\alpha, \mathrm{cod}\alpha)\}$ the natural system $B \circ (\mathrm{dom}, \mathrm{cod}) : \mathbb{C}' \to \mathrm{Ab}$. The *Hochschild-Mitchell cohomologies* $H^n(\mathbb{C}, B)$ *of the category* $\mathbb{C}$ with coefficients in the bimodule B are the Baues-Wirsching cohomologies of $\mathbb{C}$ with coefficients in the natural system $B \circ (\mathrm{dom}, \mathrm{cod})$. It is easy to see that $H^n(\mathbb{C}, B)$ isomorphic to $\lim^n_{\mathbb{C}'} \{B(\mathrm{dom}\alpha, \mathrm{cod}\alpha)\}$.

4 EXTENSION GROUPS

Let $\mathcal{A}$ be an additive category with an Abelian structure $\mathcal{P}$, and with enough $\mathcal{P}$-projectives. Then for every $A \in \mathrm{Ob}\mathcal{A}$ we have some object $P(A)$ and some proper epimorphism $\pi A : P(A) \to A$. We denote by $\omega A : \Omega(A) \to P(A)$ a kernel of πA.

Let $P_*(A)$ be the exact sequence of $\mathcal{P}$-projective objects and the proper morphisms which is obtained by sticking of sequences

$$\begin{array}{ccccccc}
0 & \to & \Omega(A) & \overset{\omega A}{\to} & P(A) & \overset{\pi A}{\to} & A & \to & 0 \\
0 & \to & \Omega^2(A) & \overset{\omega\Omega(A)}{\to} & P(\Omega(A)) & \overset{\pi\Omega(A)}{\to} & \Omega(A) & \to & 0 \\
 & & & & \cdots & & & & \\
0 & \to & \Omega^{k+1}(A) & \overset{\omega\Omega^k(A)}{\to} & P(\Omega^k(A)) & \overset{\pi\Omega^k(A)}{\to} & \Omega^k(A) & \to & 0 \\
 & & & & \cdots & & & &
\end{array}$$

That is $P_*(A)$ consists of morphisms and objects:

$$0 \leftarrow P(A) \overset{\omega A \circ \pi\Omega(A)}{\longleftarrow} P(\Omega(A)) \overset{\omega\Omega(A)\circ\pi\Omega^2(A)}{\longleftarrow} P(\Omega^2(A)) \leftarrow \dots$$
$$\leftarrow P(\Omega^k(A)) \overset{\omega\Omega^k(A)\circ\pi\Omega^{k+1}(A)}{\longleftarrow} P(\Omega^{k+1}(A)) \leftarrow \dots$$

Heller define $\mathrm{Ext}^n_{\mathcal{P}}(A,B)$ as the group $\mathrm{Hom}_{-n}(P_*(A), P_*(B))$ of homotopical classes of homogenous maps. It follows from [3, Prop. 11.6] that $\mathrm{Ext}^n_{\mathcal{P}}(A,B) \cong H^n(\mathcal{A}(P_*(A), B))$, $\forall n \geq 0$.

Denote by $L : \mathrm{Set} \to \mathrm{Ab}$ the functor from the category of sets to the category of abelian groups which assigns to each set E the free Abelian group generated by E and to each map $f : E_1 \to E_2$ the homomorphism extending f.

Lemma 4.1 *Let $\mathbb{C}$ be a small category, $\mathcal{A}$ an additive category with coproducts, $F : \mathbb{C} \to \mathcal{A}$ a functor, $D = \{D_c\}_{c \in \mathrm{Ob}\mathbb{C}}$ a family of objects $D_c \in \mathrm{Ob}\mathcal{A}$. Then $\lim^n_{\mathbb{C}'}\{\mathcal{A}(\Lambda D(\mathrm{dom}\alpha), F(\mathrm{cod}\alpha))\} = 0$, $\forall n > 0$, where $\Lambda : \mathcal{A}^{\mathrm{Ob}\mathbb{C}} \to \mathcal{A}^{\mathbb{C}}$ is the left adjoint to the restriction functor $O : \mathcal{A}^{\mathbb{C}} \to \mathcal{A}^{\mathrm{Ob}\mathbb{C}}$.*

PROOF. For every $A \in \mathrm{Ob}\mathcal{A}$ and a set E we denote by $\Sigma_E A$ the coproduct of the family $\{A_e\}_{e \in E}$ of objects $A_e = A$. Let $\nu_e : A \to \Sigma_E A$ be the canonical injections. Consider the isomorphism

$$\omega(E, A, B) : A(\Sigma_E A, B) \to \mathrm{Ab}(LE, \mathcal{A}(A, B))$$

which assigns to every morphism $\varphi : \Sigma_E A \to B$ the homomorphism $\omega(E, A, B) : LE \to \mathcal{A}(A, B)$ acting an $e \in E$ as $e \to \varphi \circ \nu_e \in \mathcal{A}(A, B)$.

Then we keep an arbitrary $c \in \mathrm{Ob}\mathbb{C}$. Consider the sets $E = h^c(a) = \mathbb{C}(c, a)$ with $a \in \mathrm{Ob}\mathbb{C}$. Let Lh^c be the composition of L and h^c. Then there is an isomorphism

$$\mathcal{A}(\Sigma_{\mathbb{C}(c,a)} A, B) \to \mathrm{Ab}(Lh^c(a), h^A(B)),$$

which is natural in $a \in \mathbb{C}$ and $A, B \in \mathcal{A}$. Let $\Lambda^c : \mathcal{A} \to \mathcal{A}^{\mathbb{C}}$ be the left adjoint to the functor $\mathcal{A}^{\mathbb{C}} \to \mathcal{A}$ acting as $F \to F(c)$ for $F \in \mathrm{Ob}(\mathcal{A}^{\mathbb{C}})$ and $\eta \to \eta_c$ for $\eta \in \mathrm{Mor}(\mathcal{A}^{\mathbb{C}})$. It is known that $(\Lambda^c A)(a) = \Sigma_{\mathbb{C}(c,a)} A$ for all $a \in \mathrm{Ob}\mathbb{C}$. For each functor $F : \mathbb{C} \to \mathcal{A}$ we have an isomorphism of bifunctors

$$\mathcal{A}(\Lambda^c A(-), F(=)) \cong \mathrm{Ab}(Lh^c(-), (h^A \circ F)(=)).$$

It leads to an isomorphism

$$\mathcal{A}(\Lambda^c A(\mathrm{dom}\alpha), F(\mathrm{cod}\alpha)) \cong \mathrm{Ab}(Lh^c(\mathrm{dom}\alpha), (h^A \circ F)(\mathrm{cod}\alpha)) \tag{3}$$

of the natural systems on $\mathbb{C}$ with fixed c. For every family $D = \{D_c\}_{c\in \mathrm{Ob}\mathbb{C}}$ of objects in $\mathcal{A}$ there exists an isomorphism $\Lambda D \cong \Sigma_{c\in\mathbb{C}}\Lambda^c D_c$. Hence

$$\lim\nolimits^n_{\mathbb{C}'}\{\mathcal{A}(\Lambda D(\mathrm{dom}\alpha), F(\mathrm{cod}\alpha))\} \cong$$

$$\lim\nolimits^n_{\mathbb{C}'}\{\textstyle\prod_{c\in\mathbb{C}} \mathcal{A}(\Lambda^c D_c(\mathrm{dom}\alpha), F(\mathrm{cod}\alpha))\} \cong$$

$$\textstyle\prod_{c\in\mathbb{C}} \lim\nolimits^n_{\mathbb{C}'}\{\mathcal{A}(\Lambda^c D_c(\mathrm{dom}\alpha), F(\mathrm{cod}\alpha))\}$$

Thus, by the isomorphism (3) it suffices to prove that

$$\lim\nolimits^n_{\mathbb{C}'}\{\mathrm{Ab}(Lh^c(\mathrm{dom}\alpha), (h^A \circ F)(\mathrm{cod}\alpha))\} = 0,$$

for every $A \in \mathrm{Ob}\mathcal{A}$ and $n > 0$. We will prove that

$$\lim\nolimits^n_{\mathbb{C}'}\{\mathrm{Ab}(Lh^c(\mathrm{dom}\alpha), G(\mathrm{cod}\alpha))\} = 0,$$

for every $c \in \mathrm{Ob}\mathbb{C}$ and $G \in \mathrm{Ab}^{\mathbb{C}}$.

For that purpose we consider the natural system

$$M = \{\mathrm{Ab}(Lh^c(\mathrm{dom}\alpha), G(\mathrm{cod}\alpha))\}$$

and the complex $K^*(\mathbb{C}, M)$. Then

$$K^n(\mathbb{C}, M) = \prod_{c_0\to\ldots\to c_n} \mathrm{Ab}(Lh^c(c_0), G(c_n)) \cong \prod_{c_0\to\ldots\to c_n}\prod_{c\to c_0} G(c_n).$$

We recall that $C^n(c/\mathbb{C}, GQ_c)$ consists of functions

$$g : N_n(c/\mathbb{C}) \to \bigcup_{\alpha\in\mathrm{Ob}(c/\mathbb{C})} GQ_c(\alpha)$$

with $g(c \to c_0 \to \ldots \to c_n) \in GQ_c(c \to c_n)$; the homomorphisms $d^n : C^n(c/\mathbb{C}, GQ_c) \to C^{n+1}(c/\mathbb{C}, GQ_c)$ act as

$$(d^n g)(c \to c_0 \to \ldots \to c_{n+1}) = \sum_{i=0}^{n}(-1)^i g(c \to c_0 \to \ldots \to \hat{c}_i \to \ldots \\ \to c_{n+1}) + (-1)^{n+1} G(c_n \to c_{n+1}) g(c \to c_0 \to \ldots \to c_n).$$

We will prove that the complex $K^*(\mathbb{C}, M)$ for

$$M = \{\mathrm{Ab}(Lh^c(\mathrm{dom}\alpha), G(\mathrm{cod}\alpha))\}$$

is isomorphic to $C^*(c/\mathbb{C}, GQ_c)$.

It is clear that $K^n(\mathbb{C}, M) \cong C^n(c/\mathbb{C}, GQ_c)$. The homomorphism $d^n : K^n(\mathbb{C}, M) \to K^{n+1}(\mathbb{C}, M)$ has the following action on functions f for which $f(c_0 \to \ldots \to c_n)(c \to c_0) \in G(c_n)$:

$$\begin{array}{ll} d^n f & (c_0 \to c_1 \to \ldots \to c_{n+1})(c \to c_0) = \\ & f(c_1 \to \ldots \to c_{n+1}) \circ Lh^c(c_0 \to c_1)(c \to c_0) + \\ & \sum_{i=1}^{n}(-1)^i f(c_0 \to \ldots \to \hat{c_i} \to \ldots \to c_{n+1})(c \to c_0) + \\ & (-1)^{n+1} F(c_n \to c_{n+1}) f(c_0 \to \ldots \to c_n)(c \to c_0). \end{array}$$

We let $\tilde{f}(c \to c_0 \to \ldots \to c_n) = f(c_0 \to \ldots \to c_n)(c \to c_0)$. We check that the correspondence $f \to \tilde{f}$ is a morphism of complexes:

$$\begin{array}{l} d^n \tilde{f}(c \to c_0 \to \ldots \to c_{n+1}) = \tilde{f}(c \to \hat{c_0} \to c_1 \to \ldots \to c_{n+1}) \\ \qquad + (-1)^i \tilde{f}(c \to c_0 \to \ldots \to \hat{c_i} \to \ldots \to c_{n+1}) + \\ \qquad (-1)^{n+1} F(c_n \to c_{n+1}) \tilde{f}(c \to c_0 \to \ldots \to c_n). \end{array}$$

Thus, the map $f \to \tilde{f}$ is an isomorphism of complexes $K^*(\mathbb{C}, M)$ and $C^*(c/\mathbb{C}, GQ_c)$. Consequently, the n-th cohomology groups of $K^*(\mathbb{C}, M)$ are zeros for $n > 0$. Hence $\lim^n_{\mathbb{C}'}\{\mathrm{Ab}(Lh^c(\mathrm{dom}\alpha), G(\mathrm{cod}\alpha))\} = 0$ for all $n > 0$. The isomorphism (3) and preservation of products by $\lim^n_{\mathbb{C}'}$ finish the proof.

Theorem 4.2 *Let $\mathcal{A}$ be an additive category with coproducts, $\mathcal{P}$ a nonempty Abelian structure. If the coproduct of every family of proper epimorphisms is proper, and $\mathcal{A}$ has enough $\mathcal{P}$-projectives, then for each small category $\mathbb{C}$ and functors $F, G : \mathbb{C} \to \mathcal{A}$ there exists a first quadrant spectral sequence with*

$$E_2^{p,q} = \lim^p_{\mathbb{C}'}\{\mathrm{Ext}^q_{\mathcal{P}}(F(\mathrm{dom}\alpha), G(\mathrm{cod}\alpha))\} \Longrightarrow \mathrm{Ext}^{p+q}_{\mathbb{C}\mathcal{P}}(F, G).$$

Proof. We will build an exact sequence

$$0 \leftarrow F \leftarrow F_0 \leftarrow F_1 \leftarrow \ldots$$

of proper morphisms with $\mathbb{C}\mathcal{P}$-projective F_n for all $n \geq 0$. Recall that $O : \mathcal{A}^{\mathbb{C}} \to \mathcal{A}^{\mathrm{Ob}\mathbb{C}}$ is the restriction functor, and Λ the left adjoint to O. The counit of adjunction $\varepsilon_F : \Lambda O F \to F$ is a retraction on each $c \in \mathrm{Ob}\mathbb{C}$. We choose a family $P = \{P(c)\}_{c \in \mathrm{Ob}\mathbb{C}}$ of $\mathcal{P}$-projectives and a family of proper epimorphisms $\psi = \{\psi_c : P(c) \to F(c)\}_{c \in \mathrm{Ob}\mathbb{C}}$ and apply the functor Λ. Then $\Lambda(\psi) : \Lambda(P) \to \Lambda O F$ is the proper epimorphism as a coproduct of proper epimorphisms. We let $F_0 = \Lambda(P)$. Let K be a kernel of the morphism $\varepsilon \circ \Lambda(\psi) : F_0 \to F$. We apply above building to K instead of F and obtain some functor K_0. We let $F_1 = K_0$. Then the inclusion $K \to F_0$ is a proper monomorphism in $\mathcal{A}^{\mathbb{C}}$. Consequently, the composition $F_1 \to K \to F_0$ is a proper morphism. We denote it by d_0. By induction we build the members $F_2, F_3, \ldots$ and morphisms $d_n : F_{n+1} \to F_n$. Now we consider the complex

$$\{K^n(\alpha)\} = \{\mathcal{A}(F_n(\mathrm{dom}\alpha), G(\mathrm{cod}\alpha))\}$$

in the category $\mathrm{Ab}^{\mathbb{C}'}$ of natural systems on $\mathbb{C}$. By Grothendieck [2] there are two spectral sequences concerned with hyper cohomologies of the functor $\lim_{\mathbb{C}'}$ with respect to the complex K^*

$$H^p(\lim^q_{\mathbb{C}'}\{K^*(\alpha)\}) \Longrightarrow H^n, \quad \lim^p_{\mathbb{C}'}\{H^qK^*(\alpha)\} \Longrightarrow H^n.$$

For each $n \geq 0$ there exists a family $A = \{A_c\}_{c\in Ob\mathbb{C}}$ of objects of $\mathcal{A}$ such that $F_n = \Lambda A$. Applying Lemma 4.1 we obtain $\lim^q_{\mathbb{C}'}\{K^n(\alpha)\} = 0$ for all $q > 0$, and $n \geq 0$. Thus, the first spectral sequence degenerates and second gives the looking spectral sequence.

The *preabelian category* is the additive category with kernels and cokernels. A morphism $f : A \to B$ in the preabelian category is called *strict* if the associated morphism $Coim\, f \to Im\, f$ is the isomorphism. Kuz'minov and Cherevikin proved that any quasiabelian category has the Abelian structure in the sense of Heller where the class of proper morphisms consists of all strict morphisms. In particular the category of locally convex spaces and coninuous linear maps is quasiabelian. Palamadov proved that this category has enough injectives. The proper monomorphisms in this category are all the kernels. This category has the infinite products. The product of kernels is the kernel. Hence, the opposite to the conditions of Theorem 4.2 are satisfied.

Corollary 4.3 *Let $\mathbb{C}$ be a small category and $\mathcal{A}$ the category of locally convex spaces and continuous linear maps. Then for every diagrams $F,\ G:\ \mathbb{C} \to \mathcal{A}$ there exists the spectral sequence with*

$$E_2^{p,q} = \lim^p_{\mathbb{C}'}\{\mathrm{Ext}^q_{\mathcal{P}}(F(\mathrm{dom}\alpha), G(\mathrm{cod}\alpha))\} \Longrightarrow \mathrm{Ext}^{p+q}_{\mathbb{C}\mathcal{P}}(F, G).$$

where $\mathcal{P}$ is the class of all strict morphisms.

References

[1] H.-J. Baues, G. Wirsching, *Cohomology of small categories.* J. Pure Appl. Algebra. 38: 2/3, (1985) 187–211.

[2] A. Grothendieck, *Sur quelques points d'algèbre homologique.* Tohoku Math. J. 9, (1957) 119–221.

[3] A. Heller, *Homological Algebra in Abelian Categories.* Ann. Math. 68: 3, (1958) 484–525.

[4] M. A. Jibladze, T. I. Pirashvili, *Some linear extensions of the category of finite generated free modules.* Soobsh. Akad. Nauk GSSR. 123: 3, (1986) 481–484. (Russian)

[5] M. Jibladze, T. Pirashvili, *Cohomology of algebraic theories.* J. Algebra. 137: 2, (1991) 253–296.

[6] A. A. Khusainov, *On extension groups in the category of Abelian diagrams.* Sib. Math. J. 33: N 1, (1992) 149–154; translation from Sib. Mat. Zh. 33: 1, (1992) 179–185.

[7] V. I. Kuz'minov, A. Yu. Cherevikin, *Semiabelian categories.* Sib. Math. J. 13, (1973) 895–902; translation from Sib. Mat. Zh. 13, (1972) 1284–1294.

[8] S. Mac Lane, *Categories for the working Mathematician.* New York, Springer-Verlag, 1971.

[9] S. Mac Lane, *Homology.* Berlin, Springer-Verlag, 1975.

[10] B. Mitchell, *Rings with several objects.* Adv. Math. 8, (1972) 1–161.

[11] Oliver, B. *Higher limits of functors on categories of elementary Abelian p-groups.* Prepr.Ser. / Mat. inst. Aarhus univ. N.6, (1992) 1–13